

The calculation of the impurity entropy for the exactly solvable two-channel Kondo model.

A.V. Rozhkov

Department of Physics, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0354, USA

In this paper the entropy of the two-channel Kondo impurity is calculated. We demonstrate that the exactly solvable system of finite size can be described in terms of free fermions. None of these fermions is Majorana fermion. This allows us to find the impurity entropy $S(T, L)$ for arbitrary values of T and L . As found before, $S(T = 0, L = \infty)$ equals to $(1/2) \ln 2$. For finite L the entropy shows a cross-over to the infinite-size behavior at $T > T_L = v_F/L$, in agreement with results of other researchers.

I. INTRODUCTION

Recent investigations of macroscopic quantum phenomenon, 1D quantum wires and Kondo effect discover that the zero-temperature entropy of some systems possesses an anomaly [1–4]. Specifically, zero-temperature entropy of the two-channel Kondo impurity was found to be:

$$S(T = 0) = \frac{1}{2} \ln 2. \quad (1)$$

This result is in obvious contradiction with the fundamental fact: for any system which exists in nature it is assumed that $\exp(S(T = 0))$ is the degeneracy of the ground state, i.e. integer number. Equation (1), however, implies that the degeneracy equals to $\sqrt{2}$.

It was pointed out later in the context of Kondo model [5,6] that the entropy value $(1/2) \ln 2$ is obtained in the thermodynamics limit where system size L is sent to infinity. For a finite size system a non-zero temperature scale can be defined in the following way: $T_L = v_F/L$. Thermodynamic limit calculations are not valid below T_L . Mathematically, this situation can be described as follows [5]:

$$\lim_{L \rightarrow \infty} \lim_{T \rightarrow 0} S = \ln 2 \neq \lim_{T \rightarrow 0} \lim_{L \rightarrow \infty} S = \frac{1}{2} \ln 2. \quad (2)$$

Only the first of these two limits is physically relevant [7] and it gives reasonable result.

As shown in [3] (further cited as I) two-channel Kondo model is exactly solvable at particular values of coupling constants. In this paper we study that exactly solvable model in the limit of finite L . A description of the finite size exactly solvable system in terms of free Dirac fermions is found. Majorana fermion which appears in $L = \infty$ system combines with another Majorana fermion and forms Dirac fermion. In $L = \infty$ limit that second Majorana fermion remains hidden in the continuous spectrum. Numerical calculation of the spectrum of the free fermions allows us to obtain the impurity entropy S for

arbitrary values of T and L . As a function of T and L the entropy demonstrates a cross-over at $T \sim T_L$, in agreement with the theory of critical phenomena. For $T > T_L$ finite size effects are unimportant while below T_L the impurity decouples from the Fermi sea. We confirm this by both numerical and analytical calculations.

The paper is organized as follows. In Section II we calculate the spectrum of the finite size version of Kondo model. As we said this model can be reduced to a set of free fermions. In Section III the thermodynamics of the model is discussed. Section IV contains the conclusions.

II. SPECTRUM OF TWO-CHANNEL KONDO IMPURITY

In this part we are going to diagonalize the hamiltonian of two-channel Kondo impurity placed in the box of size L .

In paper I two-channel Kondo hamiltonian

$$H = iv_F \sum_{i,\alpha=1}^2 \int_{-L}^L dx \psi_{i\alpha}^\dagger(x) \partial_x \psi_{i\alpha}(x) + H_I, \quad (3)$$

$$H_I = \frac{1}{2} \sum_{i,\alpha,\beta=1}^2 \sum_{\lambda=x,y,z} J_\lambda \tau_\lambda \sigma_{\alpha\beta}^\lambda \psi_{i\alpha}^\dagger(0) \psi_{i\beta}(0) \quad (4)$$

was studied in the limit of infinite L . Here ψ – left-moving electrons, α, β are electron spin indices, index i is electron "flavor", τ^λ is the impurity pseudospin. H_I describes the coupling of the electron spin to the impurity pseudospin. Coupling constants $J_{x,y,z}$ are chosen in such a way that $J_x = J_y, J_z = \pi v_F > 0$.

Following the procedure described in I (bosonization and re-fermionization) this hamiltonian can be transformed to:

$$H_{new} = iv_F \int_{-L}^L dx \Psi^\dagger(x) \partial_x \Psi(x) + \quad (5)$$

$$\frac{J_x}{\sqrt{2\pi a}} [\Psi^\dagger(0) + \Psi(0)] (d^\dagger - d),$$

where Ψ is some new fermionic field, d is a fermionic operator representing the impurity, a is a smallest length in the system (unit cell size). Expressing this hamiltonian in terms of c_k 's, Fourier components of Ψ , we obtain:

$$H_{new} = \sum_k v_F k c_k^\dagger c_k + \Delta \sum_k (c_k^\dagger + c_k)(d^\dagger - d), \quad (6)$$

$$\Delta = \frac{J_x}{\sqrt{4\pi a L}}. \quad (7)$$

Hamiltonian (5) written in terms of Fourier components (6) will be a starting point for our calculations.

To proceed further let us introduce two operators:

$$D_1 = \frac{1}{\sqrt{2}} (d^\dagger + d), \quad (8)$$

$$D_2 = \frac{i}{\sqrt{2}} (d^\dagger - d). \quad (9)$$

Obviously, D_1 and D_2 satisfy the following relations:

$$D_{1,2}^\dagger = D_{1,2}, \quad (10)$$

$$\{D_1; D_1\} = \{D_2; D_2\} = 1, \quad (11)$$

$$\{D_1; D_2\} = 0, \quad (12)$$

Though (12) is usual property of fermionic operators the equalities (10) and (11) define Majorana fermion operators. It was pointed out in I that D_1 commutes with the hamiltonian:

$$[H_{new}; D_1] = 0. \quad (13)$$

The anomalous entropy behavior in case of infinite L was attributed to the presence of this Majorana fermion commuting with (6). We will see below that there is no Majorana fermion and no anomalous entropy for the finite size system.

Now let us diagonalize the hamiltonian (6). Since it is quadratic in fermionic operators, we will look for a set of eigenoperators of the form:

$$f^\dagger = i \sum_k (\tilde{\gamma}_k c_k^\dagger + \gamma_k c_k) + \delta D_2, \quad (14)$$

such that

$$[H_{new}; f^\dagger] = \omega f^\dagger. \quad (15)$$

Conditions (14-15) give rise to an eigenvalue problem for γ 's and δ :

$$\omega \tilde{\gamma}_k = \epsilon_k \tilde{\gamma}_k - \sqrt{2} \Delta \delta, \quad (16)$$

$$\omega \gamma_k = -\epsilon_k \gamma_k - \sqrt{2} \Delta \delta, \quad (17)$$

$$\omega \delta = -\sqrt{2} \Delta \sum_k (\tilde{\gamma}_k + \gamma_k). \quad (18)$$

Here $\epsilon_k = v_F k$.

The solution of this set of equations is:

$$\tilde{\gamma}_k = -\delta \frac{\sqrt{2} \Delta}{\omega - \epsilon_k}, \quad (19)$$

$$\gamma_k = -\delta \frac{\sqrt{2} \Delta}{\omega + \epsilon_k}, \quad (20)$$

$$\omega = 2\Delta^2 \sum_k \frac{1}{\omega + \epsilon_k} + \frac{1}{\omega - \epsilon_k}. \quad (21)$$

The condition $\{f^\dagger; f\} = 1$ fixes the value of δ .

The solution shown above allows one to verify the following property of symmetry: equations (16-18) are invariant under the transformation $\omega \rightarrow -\omega, \tilde{\gamma}_k \rightarrow -\gamma_k, \gamma_k \rightarrow -\tilde{\gamma}_k, \delta \rightarrow \delta$. Another way of expressing this symmetry is:

$$f_\omega^\dagger = f_{-\omega}. \quad (22)$$

This implies that $\omega > 0$ solutions of equations (16-18) constitute fermionic spectrum of the hamiltonian (6) while $\omega < 0$ solutions are mirror image of that spectrum. In other words, $\omega > 0$ eigenvectors give us fermionic creation operators $f_\omega^\dagger$ while $\omega < 0$ eigenvectors give us destruction operators: $f_{-|\omega|}^\dagger = f_{|\omega|}$.

There is another remarkable property of equations (16-18): $\omega = 0$ solution always exists. This can be readily seen from (21). The following is true for $\omega = 0$ eigenoperator (see (22) and (15)):

$$f_{\omega=0}^\dagger = f_{\omega=0}, \quad (23)$$

$$[H_{new}; f_{\omega=0}] = 0. \quad (24)$$

Equations (23), (24) show us that $f_{\omega=0}$ is the Majorana operator (see properties (10) and (11)) commuting with the hamiltonian (6). Having two Majorana operators we can define two other operators:

$$F = \frac{1}{\sqrt{2}} (i f_{\omega=0} + D_1), \quad (25)$$

$$F^\dagger = \frac{1}{\sqrt{2}} (-i f_{\omega=0} + D_1). \quad (26)$$

These two operators are annihilation and creation operators for some new fermion:

$$\{F^\dagger; F\} = 1, \quad (27)$$

$$\{F; f_\omega\} = 0, \omega \neq 0. \quad (28)$$

Since both $f_{\omega=0}$ and D_1 commute with hamiltonian, the new fermion has zero energy:

$$[H_{new}; F] = 0. \quad (29)$$

Thus, the set of eigenoperators of H_{new} consists of f_ω for $\omega > 0$, F and their hermitian conjugates. All of them are usual fermionic operators. We have a description of our system in terms of free fermions with no Majorana fermion. The presence of the zero-energy fermion F implies that $S(T=0, L) = \ln 2$.

III. FREE ENERGY OF THE KONDO IMPURITY

In order to calculate the entropy let us consider free energy of our system. In the previous section we have shown that our system can be described in terms of free fermions. Therefore, the impurity free energy is:

$$\mathcal{F} = -T \ln 2 - T \sum_{\omega > 0} \ln \left(1 + e^{-\omega/T} \right) - \mathcal{F}_0. \quad (30)$$

where $\mathcal{F}_0$ is the free energy without the impurity. The impurity entropy is:

$$S = -\frac{\partial \mathcal{F}}{\partial T}. \quad (31)$$

We solve (21) numerically and find the entropy for different T . Three curves for three different values of L are plotted on fig.1: $L_a < L_b < L_c = \infty$. One can see from the graph that the system has a cross-over at T_L : as T goes below T_L the impurity decouples from the Fermi sea due to the presence of the finite size gaps in the excitation spectrum of the conducting electrons. It is clearly seen on fig.1 that the limiting value of $S(T, L)$ at $T = 0$ and $L = \infty$ depends on the path in (T, L) plane, in agreement with (2).




FIG. 1. The impurity entropy *vs.* temperature (in the units of Γ) for different values of L : $L_a < L_b < L_c = \infty$, $L_a/L_b = 0.6$.

There is one peculiarity on the graphs of the entropy for the finite systems. The entropy of the impurity decreases as a function of temperature below T_L . There is no contradiction here with the fundamental criterion of thermodynamical stability: the impurity entropy is not the entropy of any physical system but rather a difference between the entropy of the system with Kondo impurity and the entropy of free electrons without impurity. For both of these two systems the entropy increases with temperature, but their difference does not.

We see from fig.1 that the result for $L = \infty$ works fine as high temperature approximation. In order to show this analytically let us calculate high temperature expression for the impurity free energy (30).

In the finite size limit the spectrum consists of a set of eigenvalues defined by (21). The characteristic distance between two neighboring eigenvalues is $T_L = v_F/L$. However, for $T \gg T_L$ one can expect that discreteness of the spectrum does not play any role and it can be approximated by continuum.

In order to study continuous version of our model we are going to obtain the density of states $\rho(\omega)$. To do so we will consider the following auxiliary function

$$h(\omega) = \omega \left[\omega - 2\Delta^2 \sum_k \frac{1}{\omega + \epsilon_k} + \frac{1}{\omega - \epsilon_k} \right] \times \quad (32)$$

$$\prod_k (\omega + \epsilon_k)(\omega - \epsilon_k).$$

As it follows from (21) this function has two important properties: if ω belongs to the spectrum of our model then $h(\omega) = 0$; $h(\omega)$ does not have any singularities. It is tempting to write that $\rho(\omega) = (1/\pi)(d/d\omega)\text{Im}(\ln h(\omega))$. However, this is incorrect. There is one thing we have to take care of: for a given eigenvalue $\omega \neq 0$ the function h has two zeros (at ω and $-\omega$), for $\omega = 0$ it has a second order zero. The following is the correct expression for the density of states:

$$\rho = \frac{1}{\pi} \frac{d}{d\omega} \text{Im} \left(\frac{1}{2} \ln h(\omega) \right). \quad (33)$$

The factor of $(1/2)$ removes the impact of extra zeros of $h(\omega)$.

Then:

$$\rho = \frac{1}{2\pi} \frac{d}{d\omega} \text{Im} (\ln \omega + \quad (34)$$

$$\ln \left[\omega - 2\Delta^2 \sum_k \frac{1}{\omega + \epsilon_k} + \frac{1}{\omega - \epsilon_k} \right] + \sum_k \ln(\omega + \epsilon_k) + \sum_k \ln(\omega - \epsilon_k) \Big).$$

Expression in square brackets is evaluated in the following way:

$$\omega - 2\Delta^2 \sum_k \frac{1}{\omega + \epsilon_k} + \frac{1}{\omega - \epsilon_k} = \quad (35)$$

$$\omega - \Delta^2 \frac{2L}{\pi v_F} \int_{-\Omega}^{\Omega} \frac{d\epsilon}{\omega + \epsilon} + \frac{d\epsilon}{\omega - \epsilon} = \omega - i\Gamma,$$

where bandwidth Ω and coupling Γ are defined as follows:

$$\Omega = \frac{\pi v_F}{a}, \quad (36)$$

$$\Gamma = \frac{J_x^2}{\pi a v_F} \ll \Omega. \quad (37)$$

After straightforward calculations the density of states is found to be

$$\rho(\omega) = \rho_0 + \frac{1}{2\pi} \frac{\Gamma}{\omega^2 + \Gamma^2} + \frac{1}{2} \delta(\omega), \quad (38)$$

where ρ_0 is the density of states without the impurity.

Thus, the impurity free energy is:

$$\mathcal{F} = -T \int_{-\Omega}^{\Omega} \frac{1}{2\pi} \frac{\Gamma}{\omega^2 + \Gamma^2} \ln \left(1 + e^{-\omega/T} \right) d\omega - \frac{T}{2} \ln 2. \quad (39)$$

This free energy coincides with the free energy found in I up to constant. From (39) zero temperature value of the impurity entropy is found to be $(1/2) \ln 2$. But, as mentioned above, (39) is not valid below T_L .

IV. CONCLUSIONS

Fractional entropy paradox does not exist for the finite system. The presented calculations provide us with the direct check of that. In the case of finite system Kondo impurity can be described in terms of free fermions. It was proven above that none of them is of Majorana type. Using this representation we found the entropy S for arbitrary T and L . As expected, S shows a cross-over at $T_L = v_F/L$.

We derived high temperature expression for the impurity free energy and found that it coincides with the free energy obtained in $L = \infty$ limit.

V. ACKNOWLEDGMENTS

Author expresses his sincere gratitude to F. Guinea without whom this work could not be possible and to D. Arovas whose encouragement and advice was extremely helpful. Author would like to thank D. Cox and I. Affleck who draw his attention to the important results presented in [5].

-
- [1] F. Guinea, Phys. Rev. B **32** 7518 (1985).
 - [2] F. Lesage, H. Saleur, P. Simonetti, preprint cond-mat/9703220 v2 (1997).
 - [3] V.J. Emery, S. Kivelson, Phys. Rev. B **46** 10812 (1992).
 - [4] N. Andrei, C. Destri, Phys. Rev. Lett. **52**, 364 (1984); A.M. Tsvelik, P.B. Wiegmann, J. Stat. Phys. **38**, 125 (1985); A.M. Tsvelik, J. Phys. C **18**, 159 (1985); P.D. Sacramento, P.Schlottmann, Phys. Lett A **142**, 245 (1989).
 - [5] I. Affleck, A.W.W. Ludwig, Phys. Rev. Lett. **67**, 161 (1991); A.W.W. Ludwig, Int. J. Mod. Phys. B Vol. 8, No. 4, 1 (1994).
 - [6] P. Coleman, L. Ioffe, A. M. Tsvelik, preprint cond-mat/9504006 (1995)
 - [7] The first limit corresponds to the cooling experiment with a sample of fixed size. By cooling it low enough we have to see log of some integer. The second limit corresponds to another set of experiments when many samples of increasing size are studied at the fixed temperature. In this case there is no fundamental restriction on the limiting value of the entropy as well as that limiting value does not have any fundamental meaning.